



Pre-Admission Labs and Diagnostics

Pre-Surgery Diagnosis: _____

Scheduled Procedure: _____

Date of Procedure: _____ Allergies: _____

Precertification #: _____

No Lab Tests Required

- | | |
|---|---|
| <input type="checkbox"/> Anesthesiology Consultation (Patients with Medical or Surgical issues) | <input type="checkbox"/> MRSA Screen |
| <input type="checkbox"/> Hgb/Hct | <input type="checkbox"/> Urine HCG |
| <input type="checkbox"/> CBC without diff | <input type="checkbox"/> Serum HCG |
| <input type="checkbox"/> CBC w/diff | <input type="checkbox"/> Quant. HCG |
| <input type="checkbox"/> PT | <input type="checkbox"/> HBsAg |
| <input type="checkbox"/> PTT | <input type="checkbox"/> Sickle Cell Screening |
| <input type="checkbox"/> CMP | <input type="checkbox"/> Amylase |
| <input type="checkbox"/> BMP | <input type="checkbox"/> Lipase |
| <input type="checkbox"/> Blood Glucose | <input type="checkbox"/> Hepatic Panel |
| <input type="checkbox"/> FSBS Day of Surgery | <input type="checkbox"/> Other Lab _____ |
| <input type="checkbox"/> Uric Acid | <input type="checkbox"/> Type and Screen |
| <input type="checkbox"/> Urinalysis w/ Micro | <input type="checkbox"/> Crossmatch ___# units PRBCs |
| | <input type="checkbox"/> Crossmatch ___# units autologous Blood |
| | <input type="checkbox"/> Other blood products _____ |
| | <input type="checkbox"/> CXR |
| | <input type="checkbox"/> KUB |
| | <input type="checkbox"/> Other X-ray _____ |
| | <input type="checkbox"/> Ultrasound _____ |
| | <input type="checkbox"/> 12 lead ECG |
| | <input type="checkbox"/> PFT |

Blood for pre-transfusion testing must be drawn within 48 hours of surgery. Typed and Screened blood can only be held 48 hours.

Pre-Surgery Orders OPS (Outpatient Surgery) IP (Inpatient) Unknown Length of Stay

Vital Signs: Per Protocol Other _____

Diet: NPO NPO after _____

LR at KVO or 100ml/hr 200ml/hr _____ml/hr

NS at KVO or 100ml/hr 200ml/hr _____ml/hr

NS (500ml bag) at KVO via microdrip tubing for renal patient

Medications:

Pre-Op Antibiotic in ASU/OR holding area _____

Bacterial Endocarditis Prophylaxis: _____ Consult Pharmacy/Anesthesia

Ancef 1GM 2GM IV in OR

If PCN / B-Lactam Allergy OR Patient at Increased risk for infection

Vancomycin 1GM IV to be started within 120 minutes of incision
or

Cleocin 900MG IV to be started within 30 minutes of incision

Tylenol suppository by weight in ASU/OR

Other _____

Preparation:

Incentive Spirometry: Instructions/pre-admit

Thigh / Knee High TED hose

Sequential Compression Device in OR

Consult Anesthesia for post-op pain block

Yes No

(Type of block)

Minimum Testing Guidelines (Anesthesia Service): These guidelines are suggested minimums. They are not replacement for medical judgment, and the patient's medical history and/or the proposed surgical procedure may indicate additional laboratory or diagnostic testing. **No pre-operative laboratory testing** is required for asymptomatic patients without significant medical problems who are less than 40 years of age.

ECG: Males aged 40 and above require an ECG. Females aged 50 and above require ECG

Pregnancy Test: A pregnancy test is required for all menstruating females, unless not indicated due to sterility. Serum HCG if blood is drawn for other tests, otherwise it will be a urine HCG.

Labs: CBC and BMP are required for patients age 65 and older. BMP is required for diabetic, renal and/or hypertensive patients. FSBS is required day of surgery in diabetic patients, except in patient who have a BMP done day of surgery.

Copies: A copy of a CXR and/or ECG completed in the past 6 months is sufficient in the absence of a change in the patient's health status. A copy of lab work completed in the past 30 days is sufficient in the absence of renal disease. Renal patients must have a K+ completed after their last dialysis treatment before the date of surgery.

IVF: Every patient over age 10 is required to have an IV of LR at KVO rate. Renal patients must have NS at KVO rate