



AMBULATORY SURGERY UNIT



COFFEE REGIONAL MEDICAL CENTER	LAB & DIAGNOSTI	
Pre-Admission Labs and Diag	nostics	
Pre-Surgery Diagnosis:		
Scheduled Procedure:		
Precertification #:	-	
☐ No Lab Tests Required		
•	Patients with Medical or Sur	gical issues)
☐ Hab/Hct	☐ Urine HCG	☐ Type and Screen
☐ CBC without diff	☐ Serum Pregnancy	☐ Crossmatch# units PRBCs
	☐ Quant. HCG	Crossmatch# units autologous Blood
☐ PT	☐ HBsAg	☐ Other blood products
□ PTT	☐ Sickle Cell Screening	CXR
□ CMP	☐ Amylase	☐ KUB
□ BMP□ Blood Glucose	☐ Lipase☐ Hepatic Panel☐	☐ Other X–ray
	☐ Other Lab	☐ Ultrasound
☐ Uric Acid	- Other Lab	☐ 12 lead ECG
☐ Urinalysis w/ Micro		□ PFT
Blood for pre-transfusion testing must be drawn within 48 hours of surgery. Typed and Screened blood can only be held 48 hours.		
Pre-Surgery Orders ☐ OPS (Outpatient Surgery) ☐ IP (Inpatient) ☐ Unknown Length of Stay		
Vital Signs: Per Protocol Other	- • · · · · · · · · · · · · · · · · · ·	dient) a officiown Length of Stay
Diet: ☐ NPO ☐ NPO after		
		at KVO or 100ml/hr 200ml/hr 1ml/hr
		S at KVO or 100ml/hr 200ml/hr ml/hr
Medications:		6 (500ml bag) at KVO via microdrip tubing for renal patient
☐ Pre-Op Antibiotic in ASU/OR holding area		
□ Bacterial Endocarditis Prophylaxis:□ Consult Pharmacy/Anesthesia		
□ Ancef □ 1GM □ 2GM		
 ☐ If PCN / B-Lactam Allergy OR ☐ Patient at Increased risk for infection ☐ Vancomycin 1GM IV to be started within 120 minutes of incision 		
or	nin 120 minutes of incision	
☐ Cleocin 900MG IV to be started within	n 30 minutes of incision	
☐ Tylenol suppository by weight in ASU		Consult Anesthesia for post-op pain block
☐ Other		☐ Yes ☐ No
Preparation:		
☐ Incentive Spirometry: Instructions/pre-	-admit	
Thigh / Knee High TED hose		
Sequential Compression Device in OF	₹	(Type of block)
Minimum Testing Guidelines (Anesthesia Service): These guidelines are suggested minimums. They are not replacement for medical judgment, and the patient's medical history and/or the proposed surgical procedure may indicate additional laboratory or diagnostic testing. No pre-operative laboratory testing is required for asymptomatic patients without significant medical problems who are less than 40 years of age. ECG: Males aged 40 and above require an ECG. Females aged 50 and above require ECG		

Pregnancy Test: A pregnancy test is required for all menstruating females, unless not indicated due to sterility. Serum HCG if blood is drawn for other tests, otherwise it will be a urine HCG.

Labs: CBC and BMP are required for patients age 65 and older. BMP is required for diabetic, renal and/or hypertensive patients. FSBS is required day of surgery in diabetic patients, except in patient who have a BMP done day of surgery.

<u>Copies:</u> A copy of a CXR and/or ECG completed in the past 6 months is sufficient in the absence of a change in the patient's health status. A copy of lab work completed in the past 30 days is sufficient in the absence of renal disease. Renal patients must have a K+ completed after their last dialysis treatment before the date of surgery.

IVF: Every patient over age 10 is required to have an IV of LR at KVO rate. Renal patients must have NS at KVO rate

Name and DOB